

Modern Physics Letters A
© World Scientific Publishing Company

πK interaction effects on CP violation in $B \rightarrow K\pi^+\pi^-$ decays

B. LOISEAU

*LPNHE (IN2P3-CNRS-Univ. Paris 6 et 7), Groupe Théorie, Université Pierre et Marie Curie,
4 place Jussieu, 75252 Paris, France
loiseau@lpnhe.in2p3.fr*

B. EL-BENNICH

Physics Division, Argonne National Laboratory, Argonne, IL, 60439, USA

A. FURMAN

ul. Bronowicka 85/26, 30-091 Kraków, Poland

R. KAMIŃSKI, L. LEŚNIAK

*Division of Theoretical Physics, The Henryk Niewodniczański Institute of Nuclear Physics,
Polish Academy of Sciences, 31-342 Kraków, Poland*

B. MOUSSALLAM

IPN, CNRS/Univ. Paris-Sud 11, 91406 Orsay Cedex, France

Received (Day Month Year)

Revised (Day Month Year)

We apply QCD factorization to the quasi two-body $B \rightarrow (K\pi)\pi$ decays where the $(K\pi)$ -pair effective mass is limited to 1.8 GeV. Our strong interaction phases constrained by theory and πK experimental data yield useful information for studies of CP violation.

Keywords: B decays and QCD factorization; strange πK form factors; CP violation.

PACS Nos.: 13.25Hw, 13.75.Lb

1. Introduction

Three-body charmless non-leptonic B decays provide good opportunities for searches on CP violation in weak interactions and for studies of hadronic physics. Strong interaction phases are essential for the observation of CP violation and it is important to describe reliably the strong interaction between the detected hadrons. Here we study the three-body $B \rightarrow K\pi^+\pi^-$ decays for which BaBar and Belle Collaborations have carried out comprehensive Dalitz plot analyses^{1,2}. For $K\pi$ invariant masses, $m_{K\pi}$, below 1.8 GeV one can observe the dominance of the vector $K^*(892)$ and of the scalar $K_0^*(1430)$ resonances denoted hereafter as K^* and K_0^* .

Even if some phenomenological parameters are necessary, the Quantum Chromodynamics, QCD, factorization approach describes well B decays into two mesons³. This factorization procedure is a leading order approximation in an expansion in inverse powers of the quark b mass m_b . So far no derivation of factorization exists for B decays into three mesons. However, for $m_{K\pi} \leq 1.8$ GeV and in the rest frame of the B , the two mesons of the $K\pi$ pair move roughly in the same direction. We refer such processes as $B \rightarrow (K\pi)\pi$ and apply factorization to this quasi two-body B decay. In a preceding work the $B \rightarrow (\pi^+\pi^-)K$ decays, the final state of which is dominated by the $\rho(770)^0$ and $f^0(980)$ resonances, were studied by some of us⁴.

2. Amplitudes in the QCD factorization approach

The B -decay amplitudes are matrix elements of the effective weak Hamiltonian⁵

$$H_{eff} = \frac{G_F}{\sqrt{2}} \sum_{p=u,c} \lambda_p \left[C_1 O_1^p + C_2 O_2^p + \sum_{i=3}^{10} C_i O_i + C_{7\gamma} O_{7\gamma} + C_{8g} O_{8g} \right] + h.c. , \quad (1)$$

where, for strangeness $S = \pm 1$ final states, the λ_p are given in terms of the Cabbibo Kobayashi Maskawa, CKM, matrix elements $V_{pp'}$, $\lambda_u = V_{ub}V_{us}^*$ and $\lambda_c = V_{cb}V_{cs}^*$. Here, G_F is the Fermi coupling constant. The Wilson coefficients C_i , associated to the four-quarks operators O_i , depend on the renormalization scale μ . For the $B^+ \rightarrow (K^+\pi^-)\pi^+$ process, and in the leading order of the factorization method, one has for the penguin operators O_3 and O_4 ,

$$\begin{aligned} \langle \pi^+(K^+\pi^-) | C_3 O_3 + C_4 O_4 | B^+ \rangle = \\ a_4 \langle \pi^+ | \bar{b}\gamma^\nu (1 - \gamma_5) d | B^+ \rangle \langle K^+\pi^- | \bar{s}\gamma_\nu (1 - \gamma_5) d | 0 \rangle \end{aligned} \quad (2)$$

where $a_4 = C_4 + C_3/N_c$, N_c being the number of colors.

The short-distance physics of the weak process $\bar{b} \rightarrow \bar{s} d \bar{d}$ enters into this coefficient. Its leading order $O(\alpha_s)$ contribution $a_4(M_2)$ depends on the nature of the produced meson M_2 which, within our quasi two-body approach, we approximate to be the K_0^* for the πK - S wave or the K^* for the πK - P wave. It receives higher order QCD vertex $V_4(M_2)$ and penguin contributions^{3,5}. The penguin correction $P_4^{u(c)}(M_2)$ depends not only on M_2 but also on the intermediate quark, u or c , of the studied loop diagram. Following Refs. 5 and 6 we find, for $\mu = m_b$, $a_4 = -0.031$ for both K_0^* and K^* , and $V_4(K_0^*) = -0.001 - i0.005$, $V_4(K^*) = -0.002 - i0.001$, $P_4^u(K_0^*) = -0.029 - i0.019$ and $P_4^u(K^*) = -0.004 - i0.014$.

The transition $\langle \pi^+ | \bar{b}\gamma^\nu d | B^+ \rangle$ is given in terms of the scalar and vector B to π form factors, $F_{0,1}^{B \rightarrow \pi}(q^2)$, where the four-momentum is $q = p_B - p_{\pi^+} = p_K + p_{\pi^-}$. They contain non-perturbative physics from hadronization of quark currents^{3,5,6}. Their q^2 variation at low q^2 is small and we use $F_0^{B^+ \rightarrow \pi^+}(m_{K^*}^2) = 0.266$ and $F_1^{B^+ \rightarrow \pi^+}(m_{K^*}^2) = 0.25$. The matrix element $\langle K\pi | \bar{q}\gamma_\nu s | 0 \rangle$, given by the strange scalar and vector form factors $f_{0,1}^{K^+\pi^-}(q^2)$, have also long-range physics contributions. They are calculated⁷ using analyticity, unitarity, QCD asymptotic counting rules and accurate experimental data on πK scattering in the elastic and inelastic domains.

Adding up the next important term $a_6 = C_6 + C_5/N_c$ and neglecting other smaller contributions, the $B^+ \rightarrow (K^+\pi^-)\pi^+$ amplitude is a sum of an S - and P -wave contributions, $\mathcal{M}^+ = \mathcal{M}_S^+ + \mathcal{M}_P^+ \mathbf{p}_{\pi^-} \cdot \mathbf{p}_{\pi^+}$, with

$$\mathcal{M}_S^+ = \frac{G_F}{\sqrt{2}} (M_B^2 - m_\pi^2) \frac{m_K^2 - m_\pi^2}{q^2} F_0^{B^+ \rightarrow \pi^+}(q^2) f_0^{K^+\pi^-}(q^2) \times \left(L_4(K_0^*) - \frac{2q^2 L_6(K_0^*)}{(m_b - m_d)(m_s - m_d)} \right) \quad (3)$$

$$\mathcal{M}_P^+ = 2\sqrt{2}G_F F_1^{B^+ \rightarrow \pi^+}(q^2) f_1^{K^+\pi^-}(q^2) \left(L_4(K^*) + \frac{2m_K^*}{m_b} \frac{f_V^\perp}{f_V} L_6(K^*) \right) \quad (4)$$

where $L_i(M_2) = \lambda_u^* (a_i^u(M_2) + c_i^u) + \lambda_c^* (a_i^c(M_2) + c_i^c)$ with $a_i^p(M_2) = a_i(M_2) + V_i(M_2) + P_i^p(M_2)$ and $i = 4$ or 6 , $p = u$ or c . The term proportional to $f_V^\perp/f_V$ is inferred from the $B \rightarrow$ pseudoscalar-vector (V) amplitudes⁵. The vertex and penguin corrections are not sufficient to explain the decay rate $B \rightarrow K^*\pi$ and we have added four phenomenological complex parameters c_i^p . These could be interpreted as non perturbative contributions in the penguin diagrams.

For the $\bar{B}_0 \rightarrow (\bar{K}^0\pi^-)_{S,P} \pi^+$ amplitude one has the $b \rightarrow s\bar{u}u$ quark transition and, apart from a tree diagram a_1 contribution, the derivation is analogous to that of the $B^+ \rightarrow (K^+\pi^-)_{S,P} \pi^+$. The B^- and B_0 amplitudes are obtained from the B^+ and $\bar{B}_0$ ones with the replacements λ_u^* and λ_c^* into λ_u and λ_c .




Fig. 1. Some results from a preliminary fit to the $B \rightarrow K\pi^+\pi^-$ data. Comparison with the experimental $K\pi$ effective mass distributions for: a) $\bar{B}^0 \rightarrow \bar{K}^0\pi^+\pi^-$ and b) $B^0 \rightarrow K^0\pi^+\pi^-$. The $(K\pi)$ P -wave contribution is drawn as dotted lines, that of the S wave as dashed lines and their sum as solid lines. The data are from the Belle Collaboration².

3. Results and conclusions

We have performed a preliminary fit on the four c_i^p ($i = 4$ or 6 , $p = u$ or c). A more complete analysis will appear soon⁸. Our model reproduces well the $m_{K\pi}$

distributions in both $K^*(892)$ and $K_0^*(1430)$ regions (see Fig. 1). The S -wave contribution enhancement below 1 GeV, related to the $K^*(800)$ (or κ) resonance, is well described. The resulting $m_{\pi K}$ helicity-angle distributions compare well with experiment and show interesting S - and P -wave interference effects. The $B \rightarrow K^*(892)\pi$ branching ratios are close to the experimental values, but those for $B \rightarrow K_0^*(1430)\pi$, compatible with a new BaBar analysis, are smaller than the experimental numbers from Belle. The CP asymmetries are comparable to the experimental values.

In conclusion the use of the strange form factors⁷, constrained by experiments other than those from B decays and by theory, is an alternative to the isobar model often used in the experimental analysis of the Dalitz plots. Furthermore it allows us to calculate, using the complex pole definition of a resonance, the branching ratios and the decay constants of the $K^*(892)$ and $K_0^*(1430)$. We have looked at the effect of the next to leading order in α_s , vertex and penguin correction in the framework of the QCD factorization, it would be interesting to include also hard spectator scattering and annihilation^{3,5,6}. Nevertheless, our strong interaction phases constrained by theory and πK experimental data should provide useful information for studies of CP violation.

Acknowledgements

We acknowledge helpful comments from J-P. Dedonder and O. Leitner. This work was supported by the agreements between IN2P3 and Polish Laboratories (collaboration N° 08-127), between PAN and CNRS (collaboration N° 19481) and by the Department of Energy, Office of Nuclear Physics, contract n° DE-AC02-06CH11357.

References

1. B. Aubert, *et al.*, BaBar Collaboration, Phys. Rev. D **73**, 031101(R) (2006), *Measurements of neutral B decay branching fractions to $K_S^0\pi^+\pi^-$ final states and the charge asymmetry of $B^0 \rightarrow K^{*+}\pi^-$.*
2. A. Garmash, *et al.*, Belle Collaboration, Phys. Rev. D **75**, 012006 (2007), *Dalitz analysis of the three-body charmless $B^0 \rightarrow K^0\pi^+\pi^-$ decay.*
3. M. Beneke, Nucl. Phys. B (Proc. Suppl.) **170**, 57 (2007), *Hadronic B decays*; O. Leitner, X-H. Guo, A.W. Thomas, J. Phys. G: Nucl. Part. Phys. **31**, 199 (2005), *Direct CP violation, branching ratios and form factors $B \rightarrow \pi$, $B \rightarrow K$ in B decays.*
4. B. El-Bennich, A. Furman, R. Kamiński, L. Leśniak and B. Loiseau, Phys. Rev. D **74**, 114009 (2006), *Interference between $f_0(980)$ and $\rho(770)^0$ resonances in $B \rightarrow \pi^+\pi^-K$ decays.*
5. M. Beneke and M. Neubert, Nucl. Phys. B **675**, 333 (2003), *QCD factorization for $B \rightarrow PP$ and $B \rightarrow PV$ decays.*
6. H-Y. Cheng, C-K. Chua and K-C Yang, Phys. Rev. D **73**, 014017 (2006), *Charmless hadronic B decays involving scalar mesons: Implications to the nature of light scalar mesons.*
7. B. Moussallam, Eur. Phys. J. **C53**, 401 (2008), *Analyticity constraints on the strangeness changing vector current and applications to $\tau \rightarrow K\pi\nu_\tau$, $\tau \rightarrow K\pi\pi\nu_\tau$.*
8. B. El-Bennich, A. Furman, R. Kamiński, L. Leśniak, B. Loiseau, B. Moussallam, *CP violation and pion-kaon interactions in $B \rightarrow K\pi^+\pi^-$ decays*, work in preparation.